

**TWO NEW SPECIES OF *SEPIA* (*DORATOSEPION*) (CEPHALOPODA: SEPIIDAE) FROM  
TAIWAN, BASED ON MORPHOLOGICAL AND MOLECULAR DATA**

**Chuan-Wen Ho and Chung-Cheng Lu\***

*Department of Life Sciences, National Chung Hsing University, 250 Kuo Kuang Road,  
Taichung, Taiwan 402 (e-mail : cclu@dragon.nchu.edu.tw)*

\*corresponding author

---

ABSTRACT: Two new species of *Sepia* belonging to the *Doratosepion* group, called *Sepia furcata* and *Sepia hirunda*, are described from the waters off Taiwan, based on morphological and molecular evidence. The posterior ends of the fins of both forms extend posteriorly to form long “tails”, but there are differences in “tail” length, and the length of the arm 2 between the two species. The two species have different distributions in Taiwan: *S. furcata*, sp. nov. occurs in the northeast and *S. hirunda*, sp. nov. occurs in west and southwest of Taiwan. Analyses of genetic distance using mitochondrial DNA cytochrome *c* oxidase subunit I sequences reveal that the genetic divergence of these two species is much greater than the variation within these two species. Base on morphological and molecular data, these two taxa are considered as two valid species.

---