

POLYCHAETES OF THAILAND. NEREIDIDAE (PART 2) : *CERATOCEPHALE*
AND *GYMNONEREIS*, WITH DESCRIPTION OF TWO NEW SPECIES AND
NOTES ON THE SUBFAMILY GYMNONEREIDINAE.

by JORGEN HYLLEBERG and ANUWAT NATEEWATHANA

Phuket Marine Biological Center, P.O. Box 60, Phuket 83000, Thailand

CONTENTS

	Page
Abstract	1
I Introduction	1
II Materials and Methods	2
III Terminology	2
IV Taxonomic account	3
<i>Ceratocephale andaman</i> n.sp.	4
<i>Gymnonereis fauveli</i> (Pillai, 1961)	8
<i>Gymnonereis</i> cf. <i>fauveli</i>	10
<i>Gymnonereis phuketensis</i> n.sp.	15
<i>Gymnonereis</i> cf. <i>phuketensis</i>	17
Acknowledgements	19
References	19

ABSTRACT

A total of 150 specimens of Gymnoneridinae Banse were identified from the Andaman Sea and the Gulf of Thailand. Three species were recorded: *Ceratocephale andaman* n.sp. with and without eyes, cirrophores enlarged on setiger 9, *Gymnonereis fauveli* (Pillai, 1961) with smooth cutting edge of jaws, and *G. phuketensis* n.sp. with serrated cutting edge of jaws. The generic diagnosis of *Gymnonereis* is emended to include species with dorsal flaps and serrated jaws. The terminology of nereid parapodia is briefly reviewed and parapodia of Gymnoneridinae compared with typical Nereidinae parapodia.