

# COASTAL AND OFFSHORE PRIMARY PRODUCTION ALONG THE WEST COAST OF THAILAND (ANDAMAN SEA) WITH NOTES ON PHYSICAL-CHEMICAL VARIABLES.

BY VUDHICHAI JANEKARN and JORGEN HYLLEBERG<sup>1</sup>  
Phuket Marine Biological Center, P.O. Box 60, Phuket 83000, THAILAND.

## CONTENTS

Abstract .....	1
I. Introduction .....	1
II. Materials and methods .....	2
(a) The study area .....	2
(b) Methods .....	2
(c) Abbreviations .....	5
III. Results .....	5
(a) Physical and chemical variables .....	5
(i) Turbidity .....	5
(ii) Temperature .....	6
(iii) Salinity .....	6
(iv) Temperature-salinity relationships .....	6
(v) Nutrients .....	8
(vi) Carbonate carbon and pH .....	9
(b) Primary production and chlorophyll- <i>a</i> .....	10
(i) Northern study area from Ranong to Phuket .....	10
(ii) Southern study area from Phuket to Satun .....	15
IV. Discussion .....	19
Acknowledgements .....	20
References .....	20

## ABSTRACT

Pelagic primary production (PP) data were obtained by the <sup>14</sup>C method. *In situ* incubations were made during March-April 1982 and January-February 1983 along the west coast of Thailand in coastal and offshore areas with depths from 8 to 77 m. PP was generally high in the northern area in 1983 compared to 1982 the opposite was found in the southern area. Mean PP was 821 mgC m<sup>-2</sup> d<sup>-1</sup> (range 305–2,440) in 1982, and 888 mgC m<sup>-2</sup> d<sup>-1</sup> (range 457–1,111) in 1983. Chlorophyll-*a* concentrations were slightly higher in the northern area in 1982 than in 1983 but much higher (> 50%) in the southern area during the same years.

Physical and chemical variables were measured but no correlation could be established at this stage with PP because of complicated, interrelated processes. Temperature-salinity relationships indicated a thermocline at 40 and 45 m depth in 1982 and 1983, respectively. Upwelling is suggested to be of importance for high levels of PP measured in the study area.